

ON DIAGONALIZABLE OPERATORS IN MINKOWSKI SPACES WITH THE LIPSCHITZ PROPERTY

ZSOLT LÁNGI

ABSTRACT. A real semi-inner-product space is a real vector space $\mathbb{M}$ equipped with a function $[\cdot, \cdot] : \mathbb{M} \times \mathbb{M} \rightarrow \mathbb{R}$ which is linear in its first variable, strictly positive and satisfies the Schwartz inequality. It is well-known that the function $\|x\| = \sqrt{[x, x]}$ defines a norm on $\mathbb{M}$. and vica versa, for every norm on X there is a semi-inner-product satisfying this equality. A linear operator A on $\mathbb{M}$ is called *adjoint abelian with respect to* $[\cdot, \cdot]$, if it satisfies $[Ax, y] = [x, Ay]$ for every $x, y \in \mathbb{M}$. The aim of this paper is to characterize the diagonalizable adjoint abelian operators in finite dimensional real semi-inner-product spaces satisfying a certain smoothness condition.

1. INTRODUCTION AND PRELIMINARIES

A real semi-inner-product space is a real linear space $\mathbb{M}$ equipped with a function $[\cdot, \cdot] : \mathbb{M} \times \mathbb{M} \rightarrow \mathbb{R}$, called a *semi-inner-product*, such that

- (1) $[\cdot, \cdot]$ is linear in the first variable,
- (2) $[x, x] \geq 0$ for every $x \in \mathbb{M}$, and $[x, x] = 0$ yields that $x = 0$,
- (3) $[x, y]^2 \leq [x, x] \cdot [y, y]$ for every $x, y \in \mathbb{M}$.

These spaces were introduced in 1961 by Lumer [8], and have been extensively studied since then (cf., for example [1]). It was remarked in [8] that in a real semi-inner-product space $\mathbb{M}$, the function $\|x\| = \sqrt{[x, x]}$ defines a norm. The converse also holds, i.e. if $\mathbb{M}$ is a real linear space, then for every real norm $\|\cdot\| : \mathbb{M} \rightarrow \mathbb{R}$, there is a semi-inner-product $[\cdot, \cdot] : \mathbb{M} \times \mathbb{M} \rightarrow \mathbb{R}$ satisfying $\|x\| = \sqrt{[x, x]}$. Furthermore, the semi-inner-product determined by a norm is unique if, and only if, its unit ball is smooth; that is, if the unit sphere has a unique supporting hyperplane at its every point. By [4], in this case the semi-inner-product is homogeneous in the second variable; i.e., $[x, \lambda y] = \lambda[x, y]$ for any $x, y \in \mathbb{M}$ and $\lambda \in \mathbb{R}$.

We say that a real semi-inner-product is *continuous*, if for every $x, y, z \in \mathbb{M}$ with $[x, x] = [y, y] = [z, z] = 1$, $\lambda \rightarrow 0$ yields that $[x, y + \lambda z] \rightarrow [x, y]$ (cf. [4] or [5]). It is well-known that the semi-inner-product determined by a smooth norm is continuous; it follows, for example, from E^* on page 118 of [11] and Theorem 3 of [4].

A linear operator A is called *adjoint abelian with respect to a semi-inner-product* $[\cdot, \cdot]$, if it satisfies $[Ax, y] = [x, Ay]$ for every $x, y \in \mathbb{M}$ (cf., for instance [2] and [3]).

1991 *Mathematics Subject Classification.* 47A05, 52A21, 46B25.

Key words and phrases. semi-inner-product space, Minkowski space, norm, adjoint abelian.

In the following, $\mathbb{M}$ denotes a smooth Minkowski space; that is, a real finite dimensional smooth normed space, and $\|\cdot\|$ and $[\cdot, \cdot]$ denote the norm and the induced semi-inner-product of $\mathbb{M}$, respectively. We denote by S the unit sphere with respect to the norm, i.e., we set $S = \{x \in \mathbb{M} : \|x\| = 1\}$. We say that the semi-inner-product $[\cdot, \cdot]$ has the *Lipschitz property*, if for every $x \in S$, there is a real number κ such that for every $y, z \in S$, we have $|[x, y] - [x, z]| \leq \kappa\|y - z\|$. We note that in a similar way, a differentiability property of semi-inner-products was defined in [5], and that any semi-inner-product satisfying that differentiability property satisfies also the Lipschitz property.

The aim of this paper is to characterize the diagonalizable adjoint abelian operators in finite dimensional spaces with a semi-inner-product that satisfies the Lipschitz property.

To formulate our main result, we need the following notions and notations. An isometry of $\mathbb{M}$ is an operator $A : \mathbb{M} \rightarrow \mathbb{M}$ satisfying $\|Ax\| = \|x\|$ for every $x \in \mathbb{M}$, or, equivalently, $[Ax, Ay] = [x, y]$ for every $x, y \in \mathbb{M}$ (cf. [6]). For the properties of isometries in Minkowski spaces, the interested reader is referred to [9].

For the following definition, see also [4].

Definition 1. If $x, y \in \mathbb{M}$ and $[x, y] = 0$, we say that x is *transversal* to y , or y is *normal* to x . If $X, Y \subset \mathbb{M}$ such that $[x, y] = 0$ for every $x \in X$ and $y \in Y$, we say that X is *transversal* to Y or Y is *normal* to X .

Definition 2. Let U and V be linear subspaces of $\mathbb{M}$ such that $\mathbb{M} = U \oplus V$. If for every $x_u, y_u \in U$ and $x_v, y_v \in V$, we have

$$[x_u + x_v, y_u + y_v] = [x_u, y_u] + [x_v, y_v],$$

then we say that the semi-inner-product $[\cdot, \cdot]$ is the *direct sum* of $[\cdot, \cdot]|_U$ and $[\cdot, \cdot]|_V$, and denote it by $[\cdot, \cdot] = [\cdot, \cdot]|_U + [\cdot, \cdot]|_V$. If there are no such linear subspaces of $\mathbb{M}$, we say that $[\cdot, \cdot]$ is *non-decomposable*.

We remark that if $[\cdot, \cdot] = [\cdot, \cdot]|_U + [\cdot, \cdot]|_V$ for some linear subspaces U and V , then U and V are both transversal and normal, and that the converse does not hold. We note also that any two semi-inner-product spaces can be added in this way (cf. [5]). Definition 2 can be formulated for finitely many subspaces as well in the natural way. For the simplicity of notation, we mean that every semi-inner-product space is the direct sum of itself.

Let $A : \mathbb{M} \rightarrow \mathbb{M}$ be a linear operator, and let $\lambda_1 > \lambda_2 > \dots > \lambda_k \geq 0$ be the absolute values of the eigenvalues of A . If λ_i is an eigenvalue of A , then E_i denotes the eigenspace of A belonging to λ_i , and if λ_i is not an eigenvalue, we set $E_i = \{0\}$. We define E_{-i} similarly with $-\lambda_i$ in place of λ_i , and set $\bar{E}_i = \text{span}(E_i \cup E_{-i})$. Our main theorem is the following.

Theorem 1. Let $\mathbb{M}$ be a smooth Minkowski space such that the induced semi-inner-product $[\cdot, \cdot] : \mathbb{M} \times \mathbb{M} \rightarrow \mathbb{R}$ satisfies the Lipschitz condition, and let $A : \mathbb{M} \rightarrow \mathbb{M}$ be a diagonalizable linear operator. Then A is adjoint abelian with respect to $[\cdot, \cdot]$ if, and only if, the following hold.

- (1) $[\cdot, \cdot]$ is the direct sum of its restrictions to the subspaces $\bar{E}_i$, $i = 1, 2, \dots, k$;
- (2) for every value of i , the subspaces E_i and E_{-i} are both transversal and normal;

- (3) for every value of i , the restriction of A to $\bar{E}_i$ is the product of λ_i and an isometry of $\bar{E}_i$.

From Theorem 1, we readily obtain the following corollary.

Corollary 1. *Let $\mathbb{M}$ be a smooth Minkowski space such that the induced semi-inner-product $[\cdot, \cdot]$ satisfies the Lipschitz condition. Then the following are equivalent.*

- (1) $[\cdot, \cdot]$ is non-decomposable;
- (2) every diagonalizable adjoint abelian linear operator of $\mathbb{M}$ is a scalar multiple of an isometry of $\mathbb{M}$.

Note that if A is not diagonal, then we may apply Theorem 1 for the span of the eigenspaces of A .

Corollary 2. *Let $\mathbb{M}$ be a smooth Minkowski space such that the induced semi-inner-product $[\cdot, \cdot] : \mathbb{M} \times \mathbb{M} \rightarrow \mathbb{R}$ satisfies the Lipschitz condition, and let $A : \mathbb{M} \rightarrow \mathbb{M}$ be an adjoint abelian linear operator with respect to $[\cdot, \cdot]$. Then (1), (2) and (3) in Theorem 1 hold for A .*

If $[\cdot, \cdot] = [\cdot, \cdot]_U + [\cdot, \cdot]_V$ for some subspaces U and V , and $u \in U$ and $v \in V$, then, by Theorem 1, $S \cap \text{span}\{u, v\}$ is an ellipse. This observation is proved, for example, in Statement 1 of [5]. Thus, we have the following.

Corollary 3. *Let $\mathbb{M}$ be a smooth Minkowski space such that the induced semi-inner-product satisfies the Lipschitz condition. If no section of the unit sphere S with a plane is an ellipse with the origin as its centre, then every diagonalizable adjoint abelian operator of $\mathbb{M}$ is a scalar multiple of an isometry of $\mathbb{M}$.*

In the proof of Theorem 1, we need the following lemma.

Lemma 1. *Let $\mathbb{M}$ be a smooth Minkowski space. Let $\|\cdot\|$, $[\cdot, \cdot]$ and S denote the norm, the associated semi-inner-product and the unit sphere of $\mathbb{M}$. Then the following are equivalent.*

- (1) $[\cdot, \cdot]$ satisfies the Lipschitz condition;
- (2) for every $x \in \mathbb{M}$, the function $f_x : \mathbb{M} \rightarrow \mathbb{R}$, $f_x(y) = [x, y]$ is uniformly continuous on $\mathbb{M}$; that is, for every $x \in \mathbb{M}$ and $\varepsilon > 0$ there is a $\delta > 0$ such that $y, z \in \mathbb{M}$ and $\|y - z\| < \delta$ imply $|[x, y] - [x, z]| < \varepsilon$;
- (3) for every $x \in \mathbb{M}$ and any sequences $\{y_n\}$, $\{z_n\}$ in $\mathbb{M}$, if $\|y_n - z_n\| \rightarrow 0$, then $|[x, y_n] - [x, z_n]| \rightarrow 0$.

Proof. Note that (2) and (3) are equivalent. We prove that (1) and (3) are equivalent.

First we show that (1) yields (3). Observe that since $[x, y]$ is homogeneous in x , it suffices to prove (3) for $x \in S$. Let $x \in S$, and assume that there is a number $\kappa \in \mathbb{R}$ such that for every $y, z \in S$, we have $|[x, y] - [x, z]| < \kappa\|y - z\|$. Consider the sequences $\{y_n\}$ and $\{z_n\}$ in $\mathbb{M}$, and assume that $\|y_n - z_n\| \rightarrow 0$. Since a continuous function is uniformly continuous on any compact set and since the unit ball of $\mathbb{M}$ is compact, we may assume that $\|y_n\| \geq 1$ and that $\|z_n\| \geq 1$ for every n . Let $w_n = \frac{\|z_n\|}{\|y_n\|} y_n$. Observe that, by the definition of semi-inner-product, $|[u, v]| \leq 1$

for any $u, v \in S$. Then, from $\left\| \left[x, \frac{y_n}{\|y_n\|} \right] \right\| \leq 1$ and from the triangle inequality, we obtain that

$$\begin{aligned} |[x, y_n] - [x, z_n]| &\leq |[x, y_n] - [x, w_n]| + |[x, w_n] - [x, z_n]| = \left| \|z_n\| - \|y_n\| \right| \cdot \\ &\cdot \left\| \left[x, \frac{y_n}{\|y_n\|} \right] \right\| + \|z_n\| \cdot \left\| \left[x, \frac{w_n}{\|w_n\|} \right] - \left[x, \frac{z_n}{\|z_n\|} \right] \right\| \leq \|y_n - z_n\| + \kappa \|w_n - z_n\|. \end{aligned}$$

Note that $\|w_n - y_n\| = \left| \|y_n\| - \|z_n\| \right| \leq \|y_n - z_n\| \rightarrow 0$ and that $\|w_n - z_n\| \leq \|w_n - y_n\| + \|y_n - z_n\| \rightarrow 0$, from which it follows that $[x, y_n] - [x, z_n] \rightarrow 0$.

Assume that (1) does not hold. Then there is a point $x \in S$ and sequences $y_n, z_n \in S$ such that $[x, y_n] - [x, z_n] = \kappa_n \|y_n - z_n\|$ where $\kappa_n \rightarrow \infty$. We may assume that $\kappa_n > 0$ for every n , and since S is compact, also that $y_n \rightarrow y$ and $z_n \rightarrow z$ for some $y, z \in S$. Note that $\kappa_n \rightarrow \infty$ implies $y = z$. Let $\delta_n = \|y_n - z_n\|$, and assume that $\delta_n > 0$ for every n . Observe that as y_n and z_n converge to the same point, we have $\delta_n \rightarrow 0$, and, as $[x, y]$ is continuous in $y \in S$ for every $x \in S$, we have also that $\kappa_n \|y_n - z_n\| = \kappa_n \delta_n \rightarrow 0$. Let $u_n = \frac{y_n}{\kappa_n \delta_n}$ and $v_n = \frac{z_n}{\kappa_n \delta_n}$. Then $\|u_n - v_n\| = \frac{1}{\kappa_n} \rightarrow 0$, and $[x, u_n] - [x, v_n] = 1$, and hence, (3) does not hold. $\square$

2. PROOF OF THEOREM 1

Assume that A is adjoint abelian. Let μ and ν be two different eigenvalues of A and let x and y be eigenvectors belonging to μ and ν , respectively. Then,

$$\mu[x, y] = [Ax, y] = [x, Ay] = \nu[x, y],$$

which yields that x is transversal to y . Thus, any two eigenspaces, belonging to distinct eigenvalues, are both transversal and normal, which, in particular, proves (2) (for isometries, see this observation in [7]). Recall that an Auerbach basis of a Minkowski space is a basis in which any two distinct vectors are transversal and normal to each other, and that in every norm there is an Auerbach basis. Note that the restriction of a norm to a linear subspace is also a norm, and thus, we may choose Auerbach bases in each eigenspace separately, which, by the previous observation, form an Auerbach basis in the whole space. Let $x \in \mathbb{M}$, and observe that x has a unique representation of the form $x = \sum_{i=1}^k x_i$, where $x_i \in \bar{E}_i$. To prove Theorem 1, we need the following lemma.

Lemma 2. *If $z \in \bar{E}_i$ for some value of i , then $[z, x] = [z, x_i]$.*

Proof. Assume that $z \in \bar{E}_i$ for some $i \in \{1, 2, \dots, k\}$.

Case 1, $i = 1$.

If $\lambda_1 = 0$, then A is the zero operator, and the assertion immediately follows. Let us assume that $\lambda_1 > 0$. As A is adjoint abelian, we have that $[A^2 z, x] = [Az, Ax] = [z, A^2 x]$. Observe that $A^2 z = \lambda_1^2 z$, and that $A^2 x = \sum_{i=1}^k \lambda_i^2 x_i$. Thus,

$$(1) \quad [z, x] = \left[z, \sum_{i=1}^k \left(\frac{\lambda_i^2}{\lambda_1^2} \right)^n x_i \right]$$

for every positive integer n . Since $[\cdot, \cdot]$ is continuous in both variables, we obtain that the limit of the right-hand side of (1) is $[z, x_1]$, and hence, $[z, x] = [z, x_1]$.

Case 2, $i > 1$ and $\lambda_i \neq 0$.

We prove by induction on i . Let us assume that $[y, x] = [y, x_j]$ for every $y \in \bar{E}_j$ and $j = 1, 2, \dots, i-1$.

First, set $w = \sum_{j=1}^i x_j$ and $F_i = \text{span}(\bigcup_{j=1}^i \bar{E}_j)$. We show that $[z, w] = [z, x_i]$. Note that $A|_{F_i}$ is invertible, adjoint abelian, and its inverse is also adjoint abelian. Let B_i denote the inverse of $A|_{F_i}$, and observe that the absolute values of the eigenvalues of B_i are $\frac{1}{\lambda_j}$ and its eigenspaces are E_j and E_{-j} , where $j = 1, 2, \dots, i$. Thus, we have $\frac{1}{\lambda_i^2}[z, w] = [B_i^2 z, w] = [B_i w, B_i w] = [z, B_i^2 w]$ and

$$[z, w] = \left[z, \sum_{j=1}^i \left(\frac{\lambda_i^2}{\lambda_j^2} \right)^n x_j \right]$$

for every positive integer n . By the continuity of the semi-inner-product, and since the limit of the right-hand side is $[z, x_i]$, we have the desired equality.

Now we show that $[z, x] = [z, x_i]$. Similarly like before, we obtain that

$$[z, x] = \left[z, \sum_{j=1}^k \left(\frac{\lambda_j^2}{\lambda_i^2} \right)^n x_j \right]$$

for every positive integer n . Observe that $\lim_{n \rightarrow \infty} \left\| \sum_{j=i+1}^k \left(\frac{\lambda_j^2}{\lambda_i^2} \right)^n x_j \right\| = 0$, and that, by the previous paragraph, $\left[z, \sum_{j=1}^i \left(\frac{\lambda_j^2}{\lambda_i^2} \right)^n x_j \right] = [z, x_i]$ for every positive integer n . Thus, by Lemma 1, we have that $[z, x] = [z, x_i]$.

Case 3, $i > 1$ and $\lambda_i = 0$.

Note that in this case $i = k$ and $\bar{E}_k = E_k$. Let $F = \text{span}(\bigcup_{i=1}^{k-1} \bar{E}_i)$ and set $x_f = \sum_{j=1}^{k-1} x_j$. By Cases 1 and 2, we have $[x_f, x_k] = \sum_{j=1}^{k-1} [x_j, x_k] = 0$. On the other hand,

$$[x_k, x_f] = \left[x_k, A^2 \sum_{j=1}^{k-1} \frac{1}{\lambda_j^2} x_j \right] = \left[A^2 x_k, \sum_{j=1}^{k-1} \frac{1}{\lambda_j^2} x_j \right] = 0.$$

Thus, we obtained that F and E_k are both transversal and normal. Now we let $G = \text{span}\{x_f, x_k, z\}$.

Subcase 3.1, $\dim G = 2$.

Let $e_1 = \frac{x_f}{\|x_f\|}$ and $e_2 = \frac{x_k}{\|x_k\|}$. Since F and E_k are transversal and normal, the pair $\{e_1, e_2\}$ is an Auerbach basis in G . By Cases 1 and 2, $[\cdot, \cdot]|_F = \sum_{j=1}^{k-1} [\cdot, \cdot]|_{\bar{E}_j}$, which yields that $[e_1, \alpha_1 e_1 + \alpha_2 e_2] = \alpha_1$ for any $\alpha_1, \alpha_2 \in \mathbb{R}$.

Now we identify G with the Euclidean plane $\mathbb{R}^2$ by $\alpha_1 e_1 + \alpha_2 e_2 \mapsto (\alpha_1, \alpha_2)$; or in other words, we assume that e_1 and e_2 are the standard basis of an underlying Euclidean plane. We need to show that $[e_2, \alpha_1 e_1 + \alpha_2 e_2] = \alpha_2$ for any $\alpha_1, \alpha_2 \in \mathbb{R}$, or, equivalently, that the unit circle $S \cap G$ of the subspace G is the Euclidean unit circle.

Since $\mathbb{M}$ is smooth, $S \cap G$ is a convex differentiable curve. Consider the Descartes coordinate system induced by the standard basis e_1 and e_2 , and note that the lines $x = 1$, $x = -1$, $y = 1$ and $y = -1$ support $\text{conv } S$. Thus, for every value

of $x \in (-1, 1)$, there is exactly one point of S with x as its x -coordinate and nonnegative y -coordinate. We represent the points of $S \cap G$ with nonnegative y -coordinates as the union of the graph of a function $x \mapsto f(x)$ with $x \in [-1, 1]$, and (possibly) two segments on the lines with equations $x = 1$ and $x = -1$. We express the equality $[e_1, \alpha_1 e_1 + \alpha_2 e_2] = \alpha_1$ with the function f .

We may assume that $v = \alpha_1 e_1 + \alpha_2 e_2 \in S \cap G$. Consider the case that $v = x_0 e_1 + f(x_0) e_2$ for some $x_0 \in (-1, 1)$. Then we have $[e_1, v] = [e_1, x_0 e_1 + f(x_0) e_2] = x_0$. Let v_p denote the projection of e_1 onto the line $\{\lambda v : \lambda \in \mathbb{R}\}$ parallel to the supporting line of $\text{conv } S$ at v , and let e_p denote the projection of v onto the line $\{\lambda e_1 : \lambda \in \mathbb{R}\}$ parallel to the supporting line of $\text{conv } S$ at e_1 (cf. Figure 1). Let $v_p = \mu v$ and observe that $e_p = x_0 e_1$. We note that, by the construction of the semi-inner-product described, for example in [8], we have that $[e_1, v] = \mu$ and $[v, e_1] = x_0$. Hence the triangle with vertices o, e_1, v is similar to the triangle with vertices o, v_p, e_p , with similarity ratio x_0 . From this, we obtain that $v_p = x_0 v = x_0^2 e_1 + x_0 f(x_0) e_2$. As the line, passing through e_1 and v_p , is parallel to the supporting line of $\text{conv } S$ at v , we have

$$f'(x_0) = -\frac{x_0 f(x_0)}{1 - x_0^2},$$

which is an ordinary differential equation for f with the initial condition $f(0) = 1$.




FIGURE 1. An illustration for Subcase 3.1

We omit an elementary computation that shows that the solution of this differential equation is $y = \sqrt{1 - x^2}$. Thus, we obtain that $S \cap G$ is the Euclidean unit circle, which yields, in particular, that $[z, x_f + x_k] = [z, x_k]$.

Subcase 3.2, $\dim G = 3$.

Set $e_1 = \frac{x_f}{\|x_f\|}$ and choose an Auerbach basis $\{e_2, e_3\}$ in $\text{span}\{x_k, z\}$. Then the set $\{e_1, e_2, e_3\}$ is an Auerbach basis in G . Furthermore, since F and E_k are transversal and normal, $\{e_1, v\}$ is an Auerbach basis in its span for any $v \in \text{span}\{x_k, z\}$ with $\|v\| = 1$. Thus, applying the argument in Subcase 3.1 for the subspace $\text{span}\{e_1, v\}$, we obtain that $S \cap \text{span}\{e_1, v\}$ is the ellipse with semi-axes e_1 and v . Note that this property and $S \cap \text{span}\{e_1, e_2\}$ determines the norm.

Consider the semi-inner-product defined by $[\beta_1 e_1 + \beta_2 e_2 + \beta_3 e_3, \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3]' = \beta_1 \alpha_1 + [\beta_2 e_2 + \beta_3 e_3, \alpha_2 e_2 + \alpha_3 e_3]$. We show that $[\cdot, \cdot]'$ and $[\cdot, \cdot]$ define the same norm, which, as a smooth norm uniquely determines its semi-inner-product, yields that $[\cdot, \cdot]' = [\cdot, \cdot]$.

Let $v = \alpha_2 e_2 + \alpha_3 e_3$ with $\|v\| = 1$ be arbitrary. Note that if $\|\mu e_1 + \nu v\| = 1$, then

$$[\mu e_1 + \nu v, \mu e_1 + \nu v]' = \mu^2 + \nu^2 [v, v] = \mu^2 + \nu^2 = 1,$$

which, in $\text{span}\{e_1, v\}$, is the equation of the ellipse with semiaxes e_1 and v . As the restrictions of $[\cdot, \cdot]'$ and $[\cdot, \cdot]$ to $\text{span}\{e_2, e_3\}$ are clearly equal, we obtain that $[\cdot, \cdot] = [\cdot, \cdot]'$, which, in particular, implies that $[z, x_f + x_k] = [z, x_k]$. $\square$

By Lemma 2, we have that (1) of Theorem 1 holds. Thus, it remains to show that (3) also holds. Without loss of generality, let us assume that $k = 1$, and that $\lambda_1 = 1$. Then every $x \in \mathbb{M}$ can be decomposed as $x = x_1 + y_1$ with $x_1 \in E_1$ and $y_1 \in E_{-1}$. Hence,

$$[A(x_1 + y_1), A(x_1 + y_1)] = [x_1 + y_1, A^2(x_1 + y_1)] = [x_1 + y_1, x_1 + y_1],$$

and thus, A is an isometry.

Finally, we show that if (1), (2) and (3) holds, then A is adjoint abelian. Let $x = \sum_{i=1}^k x_i$ and $y = \sum_{i=1}^k y_i$ with $x_i, y_i \in \bar{E}_i$. Assume, first, that $\lambda_k \neq 0$, which means that A is invertible. Note that $\bar{E}_i$ is an invariant subspace of A for every value of i , and that $\left(\frac{1}{\lambda_i} A\right)^{-1} = \frac{1}{\lambda_i} A = \text{id}$ on $\bar{E}_i$. Hence,

$$\begin{aligned} [Ax, y] &= \sum_{i=1}^k [Ax_i, y_i] = \sum_{i=1}^k \lambda_i \left[\frac{1}{\lambda_i} Ax_i, y_i \right] = \sum_{i=1}^k \lambda_i \left[x_i, \left(\frac{1}{\lambda_i} A \right)^{-1} y_i \right] = \\ &= \sum_{i=1}^k \lambda_i \left[x_i, \frac{1}{\lambda_i} Ay_i \right] = \sum_{i=1}^k [x_i, Ay_i] = [x, Ay], \end{aligned}$$

and the assertion follows. If A is not invertible, we may apply a slightly modified argument.

Acknowledgements. The author is indebted to Á. G. Horváth for proposing this project and for his helpful comments.

REFERENCES

- [1] S. S. Dragomir, *Semi-inner-products and Applications* Nova Science Publishers, Hauppauge NY, 2004.
- [2] R. J. Fleming and J. E. Jamison, *Hermitian and adjoint abelian operators on certain Banach spaces*, Pacific J. Math. **52** (1974), 67-84.
- [3] R. J. Fleming and J. E. Jamison, *Adjoint abelian operators on L^p and $C(K)$* , Trans. Amer. Math. Soc. **217** (1976), 87-98.
- [4] J. R. Giles, *Classes of semi-inner product spaces*, Trans. Amer. Math. Soc. **129** (1967), 436-446.
- [5] Á. G. Horváth, *Semi-indefinite product and generalized Minkowski spaces*, J. Geom. Phys. **60** (2010), 1190-1208.
- [6] D. O. Koehler, *A note on some operator theory in certain semi-inner-product spaces*, Proc. Amer. Math. Soc. **30**(2) (1971), 363-366.
- [7] D. Kohler and P. Rosenthal, *On isometries of normed linear spaces*, Studia Math. **36** (1970), 213-216.
- [8] G. Lumer, *Semi-inner-product spaces*, Trans. Amer. Math. Soc. **100** (1961), 29-43.

- [9] H. Martini and M. Spirova, *Reflections in strictly convex Minkowski planes*, Aequationes Math. **78** (2009), 71-85.
- [10] R. E. Megginson, *An Introduction to Banach Space Theory*, Graduate texts in mathematics **183**, Springer-Verlag, New York, 1998.
- [11] A. W. Roberts, D. E. Varberg, *Convex functions*, Pure and Applied Mathematics **57**, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1973.

ZSOLT LÁNGI, DEPT. OF GEOMETRY, BUDAPEST UNIVERSITY OF TECHNOLOGY, BUDAPEST,
EGRY JÓZSEF U. 1., HUNGARY, 1111

E-mail address: `zlangi@math.bme.hu`